

7 A1 a) $z^3 - 6z^2 + 13z = 0$

$z \cdot (z^2 - 6z + 13) = 0 \quad z_1 = 0$

$z^2 - 6z + 13 = 0 \quad D = (-6)^2 - 4 \cdot 13 = 36 - 52 = -16$

$z_{2,3} = \frac{6 \pm \sqrt{-16}}{2} = \frac{6 \pm 4i}{2} = 3 \pm 2i$

$z_2 = 3 + 2i \quad z_3 = 3 - 2i$

4 b) $z^4 - 8z^2 - 9 = 0$

Sub.: $z^2 = u$

$u^2 - 8u - 9 = 0$

Vieta: $(u-9) \cdot (u+1) = 0 \quad u_1 = 9 \quad u_2 = -1$

Result: $z^2 = 9 \quad z_1 = 3 \quad z_2 = -3$

$z^2 = -1 \quad z_3 = i \quad z_4 = -i$

4 A2 $z_1 = 1 + 2i \Rightarrow z_2 = \overline{z_1} = 1 - 2i$

$(z - z_1) \cdot (z - z_2) = (z - (1 + 2i)) \cdot (z - (1 - 2i))$

$= z^2 - 2z + 5$

Polynom division: $(z^4 - 2z^3 + 3z^2 + 4z - 10) : (z^2 - 2z + 5) =$

$$\begin{array}{r} z^2 - 2 \\ \underline{z^4 - 2z^3 + 3z^2 + 4z - 10} \\ z^4 - 2z^3 + 5z^2 \\ \underline{ - 2z^2 + 4z - 10} \\ - 2z^2 + 4z - 10 \\ \underline{ 0} \end{array}$$

$z^2 - 2 = 0$

$z^2 = 2 \Rightarrow z_3 = \sqrt{2} \quad ; \quad z_4 = -\sqrt{2}$

3 A3 a) z.B $f(x) = x \cdot (x-1) \cdot (x^2+1)$

b) z.B $g(x) = (x^2+1)^2$

c) z.B $h(x) = x^2 \cdot (x^2+1) \cdot (x^2+4)$

3 A4 $\frac{2x+6}{x^2-1} = \frac{2x+6}{(x-1) \cdot (x+1)} = \frac{A}{x-1} + \frac{B}{x+1}$

$A(x+1) + B(x-1) = \underbrace{(A+B)}_2 x + \underbrace{A-B}_6$

$B = 2 - A = 2 - 4 = -2$

$\begin{array}{l} A+B=2 \\ A-B=6 \quad | + \\ \hline 2A=8 \quad A=4 \end{array}$

$$A4 \quad \frac{2x+6}{x^2-1} = \frac{4}{x-1} + \frac{-2}{x+1} = \frac{4}{x-1} - \frac{2}{x+1} - 2 -$$

$$14A5_4 a) \int_0^1 \underbrace{4x}_{u} \cdot \underbrace{e^{-0,5x}}_{v'} dx = \int \quad u' = 4 \quad v = -2e^{-0,5x}$$

$$\int = \left[-8x e^{-0,5x} \right]_0^1 - \int_0^1 -8 e^{-0,5x} dx$$

$$\int = -8e^{-0,5} + \left[-16e^{-0,5x} \right]_0^1$$

$$\int = -\frac{8}{\sqrt{e}} - \frac{16}{\sqrt{e}} - (-16) = 16 - \frac{24}{\sqrt{e}}$$

$$5 b) \quad \frac{15}{(x-1) \cdot (x+2)} = \frac{A}{x-1} + \frac{B}{x+2}$$

$$A \cdot (x+2) + B(x-1) = 15$$

$$\underbrace{(A+B)}_0 x + \underbrace{2A-B}_{15} = 15 \quad \text{für alle } x \in \mathbb{R} \setminus \{1, -2\}$$

$$\begin{array}{l} A+B=0 \\ 2A-B=15 \end{array} \quad + \Rightarrow \quad \begin{array}{l} 3A=15 \\ 3B=-15 \end{array} \Rightarrow \begin{array}{l} A=5 \\ B=-5 \end{array}$$

$$\int_2^3 \frac{15}{(x-1) \cdot (x+2)} dx = \int_2^3 \frac{5}{x-1} dx - \int_2^3 \frac{5}{x+2} dx$$

$$= \left[5 \ln(x-1) \right]_2^3 - \left[5 \ln(x+2) \right]_2^3$$

$$= 5 \ln 2 - 5 \underbrace{\ln 1}_0 - (5 \ln 5 - 5 \ln 4)$$

$$= 5 \cdot (\ln 2 - \ln 5 + \ln 4) = 5 \cdot \ln \frac{8}{5}$$

Innere
Ableitung
↑
des
Wurzel

$$5 c) \int_0^3 \frac{4x}{\sqrt{25-x^2}} dx = \int_0^3 (-2) \cdot \frac{-2x}{\sqrt{25-x^2}} dx = -2 \cdot \int_0^3 \frac{-2x}{\sqrt{25-x^2}} dx$$

$$= -2 \left[2 \cdot (25-x^2)^{\frac{1}{2}} \right]_0^3 = -2 \cdot (2 \cdot \sqrt{16} - 2 \cdot \sqrt{25})$$

$$= -2 \cdot (8 - 10) = -2 \cdot (-2) = 4$$

$$5 A6 \text{ Substitution: } x = e^u \Rightarrow \frac{dx}{du} = e^u \Rightarrow dx = e^u \cdot du$$

$$\text{Neue Grenze: } e^u = 1 \Rightarrow u_1 = 0 \quad ; \quad e^u = 2 \Rightarrow u_2 = \ln 2$$

$$\int_1^2 \frac{\ln(x^2)}{x} dx = \int_0^{\ln 2} \frac{\ln((e^u)^2)}{e^u} \cdot e^u du = \int_0^{\ln 2} \ln(e^{2u}) du$$

$$= \int_0^{\ln 2} 2u du = \left[u^2 \right]_0^{\ln 2} = (\ln 2)^2$$

19VP

$$4 A7_1 a) A^T = \begin{pmatrix} 2 & 0 & 1 & -2 \\ -1 & 1 & 2 & 2 \\ 0 & 3 & -3 & 1 \end{pmatrix}$$

-3-

$$2 b) A \cdot B = \begin{pmatrix} 2 & -1 & 0 \\ 0 & 1 & 3 \\ 1 & 2 & -3 \\ -2 & 2 & 1 \end{pmatrix} \cdot \begin{pmatrix} 3 & 1 \\ -1 & 2 \\ 0 & 3 \end{pmatrix} = \begin{pmatrix} 7 & 0 \\ -1 & 11 \\ 1 & -4 \\ -8 & 5 \end{pmatrix}$$

1 c) Da B zwei Spalten und A vier Zeilen hat.
Es müsste jedoch die gleiche Anzahl sein, damit die Multiplikation definiert ist.

$$2 A8 \begin{array}{ccc} \cancel{1} & \cancel{4} & \cancel{0} \\ \cancel{-2} & \cancel{0} & \cancel{-4} \\ \cancel{3} & \cancel{-1} & \cancel{2} \\ \cancel{1} & \cancel{4} & \cancel{0} \\ \cancel{-2} & \cancel{0} & \cancel{-4} \end{array} \quad \det(C) = 0 + 0 + (-48) - 0 - 4 - (-16)$$

$$\det(C) = -36$$

$$4 A9 \text{ Ansatz } \begin{pmatrix} -1 & 1 \\ 4 & 2 \end{pmatrix} \cdot \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow \begin{array}{l} -a + c = 1 \\ 4a + 2c = 0 \end{array} \begin{array}{l} \cdot 4 \\ + \end{array} \quad \text{bzw.} \quad \begin{array}{l} -b + d = 0 \\ 4b + 2d = 1 \end{array} \begin{array}{l} \cdot 4 \\ + \end{array}$$

$$6c = 4$$

$$\Rightarrow c = \frac{2}{3} \Rightarrow a = -\frac{1}{3}$$

$$6d = 1$$

$$\Rightarrow d = \frac{1}{6} \Rightarrow b = \frac{1}{6}$$

$$D^{-1} = \begin{pmatrix} -\frac{1}{3} & \frac{1}{6} \\ \frac{2}{3} & \frac{1}{6} \end{pmatrix}$$

8VP

$\Sigma 46VP$

Alternative Lösung zu 5c)

$$\text{Substitution: } x = 5 \sin u \Rightarrow \frac{dx}{du} = 5 \cos u \Rightarrow dx = 5 \cos u \cdot du$$

$$\text{Neue Grenze: } 5 \sin u = 0 \Rightarrow u_1 = 0; \quad 5 \sin u = 3 \Rightarrow u_2 = \arcsin \frac{3}{5}$$

$$\int_0^{\arcsin 0,6} \frac{4x}{\sqrt{25-x^2}} dx = \int_0^{\arcsin 0,6} \frac{4 \cdot 5 \sin u}{\sqrt{25-25 \sin^2 u}} \cdot 5 \cos u du$$

$$= \int_0^{\arcsin 0,6} \frac{20 \sin u \cdot 5 \cos u}{\sqrt{25 \cdot (1-\sin^2 u)}} du = \int_0^{\arcsin 0,6} \frac{20 \sin u \cdot 5 \cos u}{5 \cdot \cos u} du = \int_0^{\arcsin 0,6} 20 \sin u du$$

$$= \left[-20 \cos u \right]_0^{\arcsin 0,6} = -20 \cdot (\cos(\arcsin 0,6) - 1)$$

$$= -20 \cdot (\sqrt{1-\sin^2(\arcsin 0,6)} - 1) = -20 \cdot (\sqrt{1-0,6^2} - 1)$$

$$= -20 \cdot (\sqrt{0,64} - 1) = -20 \cdot (0,8 - 1) = -20 \cdot (-0,2) = 4$$